arXiv:1406.5579v1 [math.RA] 21 Jun 2014

Some Results about Triangular Representations of Lie Algebras

Keqin Liu
Department of Mathematics
The University of British Columbia
Vancouver, BC
Canada, V6T 1Z2

June , 2014

Abstract

We introduce the concept of a triangular representation of a Lie algebra, give a counterpart of Ado's theorem, and discuss 2-irreducible triangular modules over a nonreductive Lie algebra.

It is well-known that the representation theory of semisimple Lie algebras is a fruitful area in Lie theory, but it is basically not known how to develop the finite dimensional representation theory of non semisimple Lie algebras. This fact makes us feel strongly that the ordinary definition of a representation of a Lie algebra is too general to give rise to an effective way of developing the representation theory of non semisimple Lie algebras. If a restrictive definition has to be used to study representations for non semisimple Lie algebras, which conditions should be added to the ordinary definition of a representation of a Lie algebra?

By a fact which is due to E. Cartan, a nonreductive Lie algebra does not have a faithful finite dimensional irreducible representation. Hence, replacing the linear transformations in the general linear Lie algebra by the linear transformations which have more than two invariant subspaces should be one change we should do to introduce a restrictive definition for a representation of a non semisimple Lie algebra. By Ado's theorem, a finite dimensional Lie algebra over a field of characteristic 0 has a finite dimensional faithful representation such that every element in its nil radical (which is the largest nilpotent ideal) can be represented by a nilpotent linear transformation. Since it is easy to get a lot of nilpotent linear transformations on a finite dimensional graded vector space and a finite dimensional module over a finite dimensional Lie algebra automatically has a graded vector space structure after treating the module as a module over a Levi factor of the Lie algebra, replacing the general vector space in the ordinary definition of a representation of a Lie algebra by a graded vector space and

requiring that the elements in the nil radical of a Lie algebra are represented in terms of a class of obvious nilpotent linear transformations on the graded vector space should be another change we should do. Based on these ideals, we write this paper to introduce the concept of a triangular representation of a Lie algebra and to give some evidences which indicate that it is reasonable to develop the representation theory of non semisimple Lie algebras by using triangular representations.

This paper consists of two parts. In part 1, we use linear transformations which have many invariant subspaces to define the general triangular linear Lie algebra. It turns out that the general triangular linear Lie algebra, which is a subalgebra of the general linear Lie algebra, has a natural graded vector space structure. Based on this property of the general triangular linear Lie algebra, we add two more conditions in the ordinary definition of a representation of a Lie algebra to introduce the concept of a triangular representation of a Lie algebra. We finish Part 1 with the main result of this paper which claims that every finite dimensional Lie algebra over a field of characteristic 0 has a finite dimensional faithful triangular representation. Because of this counterpart of Ado's theorem, we know that although the concept of a triangular representation of a Lie algebra is restrictive, this concept is still general enough. This is our reason to initiate the study of the triangular representations of Lie algebras. In Part 2, we discuss finite dimensional 2-irreducible triangular representations of a nonreductive Lie algebra $s\ell_2^\Lambda$, where $s\ell_2^\Lambda$ is the semidirect product of the 3-dimensional Lie algebra $s\ell_2$ by the radical $rad(s\ell_2^\Lambda)$ of $s\ell_2^\Lambda$ such that $rad(s\ell_2^\Lambda)$ is both an Abelian subalgebra of $s\ell_2^\Lambda$ and a finite dimensional irreducible $s\ell_2$-module.

Throughout this paper, $\mathbf{F}$ is a field of characteristic 0, all vector spaces are finite dimensional vector spaces over $\mathbf{F}$, and the notation $k \in \mathbf{Z}_{\geq j}$ means that k is an integer which is larger than or equal to j.

Part 1: The counterpart of Ado's theorem

For convenience, we use $\bigoplus_{k\geq j} V_k$ to denote a graded vector space $\bigoplus_{k\in\mathbf{Z}_{\geq j}} V_k$. If $V = \bigoplus_{k\geq j} V_k$ is a finite dimensional graded vector space, then $V_k = \{0\}$ for large k.

Definition 0.1 *Let $V = \bigoplus_{k\geq 0} V_k$ be a finite dimensional graded vector space. A* **triangular linear transformation** *$f : V \to V$ is a linear transformation on V satisfying the condition: $f\big(\bigoplus_{k\geq j} V_k\big) \subseteq \bigoplus_{k\geq j} V_k$ for $j \in \mathbf{Z}_{\geq 0}$.*

Let $End^{\triangle}\big(\bigoplus_{k\geq 0} V_k\big)$ be the set of all triangular linear transformations on the graded vector space $\bigoplus_{k\geq 0} V_k$. We write $g\ell^{\triangle}\big(\bigoplus_{k\geq 0} V_k\big)$ for $End^{\triangle}\big(\bigoplus_{k\geq 0} V_k\big)$ viewed as a subalgebra of the general linear Lie algebra $g\ell\big(\bigoplus_{k\geq 0} V_k\big)$ and call it the **general triangular linear Lie algebra**.

Proposition 0.1 *If $V = \bigoplus_{k\geq 0} V_k$ is a finite dimensional graded vector space, then $g\ell^{\triangle}(V) = \bigoplus_{j\geq 0} g\ell_j^{\triangle}(V)$ as the direct sum of vector spaces, where $g\ell_j^{\triangle}(V)$ with $j \in \mathbf{Z}_{\geq 0}$ is defined by*

$$g\ell_j^{\triangle}(V) := \{\, f \,|\, f \in g\ell^{\triangle}(V) \text{ and } f(V_k) \subseteq V_{k+j} \text{ for } k \in \mathbf{Z}_{\geq 0} \,\}.$$

Obviously, every element in $\bigoplus_{j\geq 1} g\ell_j^{\triangle}(V)$ is a nilpotent linear transformation. Using these nilpotent linear transformations in $\bigoplus_{j\geq 1} g\ell_j^{\triangle}(V)$, we now introduce the concept of a triangular representation of a Lie algebra.

Definition 0.2 *Let $\mathcal{L}$ be a Lie algebra and let $V = \bigoplus_{k\geq 0} V_k$ be a finite dimensional graded vector space. A* **triangular representation** *of the Lie algebra $\mathcal{L}$ is a Lie algebra homomorphism $\varphi : \mathcal{L} \to g\ell^{\triangle}(V)$ which satisfies the following two more conditions:*

(i) *$\varphi(\mathcal{L}_{semi}) \subseteq g\ell_0^{\triangle}(V)$, where $\mathcal{L}_{semi}$ is a Levi factor of $\mathcal{L}$;*

(ii) *$\varphi(nil\,\mathcal{L}) \subseteq \bigoplus_{j\geq 1} g\ell_j^{\triangle}(V)$, where $nil\,\mathcal{L}$ is the nil radical of $\mathcal{L}$.*

The condition (i) in Definition 0.2 makes it possible to use the fruitful representation theory of semisimple Lie algebras in the study of triangular representations of non semisimple Lie algebras. Because the condition (i) in Definition 0.2 depends on a Levi factor of a Lie algebra, a triangular representation $\varphi : \mathcal{L} \to g\ell^{\triangle}(V)$ is also called the triangular representation of $\mathcal{L}$ with respect to the Levi factor $\mathcal{L}_{semi}$, and the graded vector space V is called the **triangular $\mathcal{L}$-module** with respect to the Levi factor $\mathcal{L}_{semi}$. By the following proposition, the concept of triangular representation of a Lie algebra does not depends on the choice of a Levi factor of the Lie algebra.

Proposition 0.2 *Let $\mathcal{L}$ be a finite dimensional Lie algebra. If a finite dimensional graded vector space $V = \bigoplus_{k\geq 0} V_k$ is a triangular $\mathcal{L}$-module with respect to a Levi factor of the Lie algebra $\mathcal{L}$, then V is also a triangular $\mathcal{L}$-module with respect to any Levi factor of the Lie algebra $\mathcal{L}$.*

The next proposition proves that the concept of triangular representation of a Lie algebra is in fact quite natural.

Proposition 0.3 *A finite dimensional Lie algebra can be made into a graded vector space such that its adjoint representation is a triangular representation.*

We finish Part 1 of this paper with the following counterpart of Ado's theorem.

Proposition 0.4 *A finite dimensional Lie algebra over a field of characteristic 0 has a finite dimensional faithful triangular representation.*

Part 2: Triangular 2-irreducible $s\ell_2^\Lambda$-modules

Let $\mathcal{L}$ be a finite dimensional Lie algebra and $n \in \mathbf{Z}_{\geq 0}$. A finite dimensional graded vector space $V = \bigoplus_{k=0}^{n} V_k$ with respect to a Levi factor $\mathcal{L}_{semi}$ of $\mathcal{L}$ is said to be $(n+1)$**-irreducible** if each V_k is an irreducible $\mathcal{L}_{semi}$-module for $k = 0$, 1, ... , n.

The purpose of Part 2 of this paper is to describe 2-irreducible triangular $s\ell_2^\Lambda$-modules, where the nonreductive Lie algebra $s\ell_2^\Lambda$ with $\Lambda \in \mathbf{Z}_{\geq 1}$ is defined as follows:

$$s\ell_2^\Lambda := \mathbf{F}f \oplus \mathbf{F}h \oplus \mathbf{F}e \oplus \mathbf{F}z_0 \oplus \mathbf{F}z_1 \oplus \cdots \oplus \mathbf{F}z_\Lambda,$$

$$[h,\, e] = 2e, \qquad [h,\, f] = -2f, \qquad [e,\, f] = h, \qquad [z_j,\, z_{j'}] = 0,$$

$$[h,\, z_j] = (\Lambda - 2j)z_j, \qquad [f,\, z_j] = z_{j+1}, \qquad [e,\, z_j] = j(\Lambda - j + 1)z_{j-1},$$

where $\Lambda \geq j,\, j' \geq 0$ and $z_j := 0$ for $j \notin \{0,\, 1,\, ..., \Lambda\}$.

Clearly, the 3-dimensional Lie algebra $s\ell_2 = \mathbf{F}f \oplus \mathbf{F}h \oplus \mathbf{F}e$ is a Levi factor of $s\ell_2^\Lambda$, and $\mathbf{F}z_0 \oplus \mathbf{F}z_1 \oplus \cdots \oplus \mathbf{F}z_\Lambda$ is both the radical and the nil radical of $s\ell_2^\Lambda$.

For arbitrary four nonnegative integers m, n, s and N which satisfy the conditions:

$$m + 2s = \Lambda + n + 2N, \qquad n \geq s \geq 0, \qquad m \geq N \geq 0, \tag{1}$$

and arbitrary $n - s$ scalars a_1, ... , $a_{n-s} \in \mathbf{F}$, there is a 2-irreducible triangular $s\ell_2^\Lambda$-module $\mathcal{M}_{m,\,n}^{s,\,N}(a_1, ..., a_{n-s})$ which is defined as follows:

$$\mathcal{M}_{m,\,n}^{s,\,N}(a_1, ..., a_{n-s}) = \Big(\bigoplus_{i=0}^{n} \mathbf{F}u_i\Big) \bigoplus \Big(\bigoplus_{k=0}^{m} \mathbf{F}w_k\Big);$$

$$h \cdot u_i = (n-2i)u_i, \quad f \cdot u_i = u_{i+1}, \quad e \cdot u_i = i(n-i+1)u_{i-1} \quad \text{for } n \geq i \geq 0;$$

$$h \cdot w_k = (m-2k)w_k, \quad f \cdot w_k = w_{k+1}, \quad e \cdot w_k = k(n-k+1)w_{k-1} \quad \text{for } m \geq k \geq 0;$$

$$z_j \cdot w_k = 0 \quad \text{for } \Lambda \geq j \geq 0 \text{ and } m \geq k \geq 0;$$

$$z_j \cdot v_i = 0 \quad \text{for } \Lambda \geq j \geq 0,\, n \geq i \geq 0 \text{ and } s - 1 \geq i + j;$$

$$z_j \cdot v_{s-j+\theta} = \left(\sum_{k=0}^{\theta} (-1)^{j-k} \binom{j}{k} \frac{(m-N-\theta+k)!}{(N+\theta-k)!\, m!}\, a_{\theta-k}\right) w_{\theta+N}$$

$$\text{for } min\{\, s+\theta,\, \lambda \,\} \geq j \geq 0 \text{ and } n - s \geq \theta \geq 0;$$

$$z_j \cdot v_{n-j+\theta} = \left(\sum_{k=0}^{j-\theta} (-1)^{j-\theta-k} \binom{j}{\theta+k} \frac{(m-N-n+s+k)!}{(N+n-s-k)!\, m!} a_{n-s-k}\right) w_{n-s+\theta+N}$$

for $\lambda \geq j \geq \theta \geq 1$,

where $a_0 := 1$, $\binom{j}{k} := \frac{j!}{k!(j-k)!}$ for $j \geq k$, $\binom{j}{k} := 0$ for $k > j$, $u_i := 0$ for $i \notin \{0, 1, ..., n\}$, and $w_k := 0$ for $k \notin \{0, 1, ..., m\}$.

We finish this paper with the following

Proposition 0.5 *Let $\mathcal{M}$ be a 2-irreducible triangular module over the non-reductive Lie algebra $s\ell_2^\Lambda$. If $rad(s\ell_2^\Lambda) \cdot \mathcal{M} \neq \{0\}$, then $\mathcal{M} = \mathcal{M}_{m,n}^{s,N}(a_1, ..., a_{n-s})$ for some $n - s$ scalars a_1, ... , $a_{n-s} \in \mathbf{F}$ and some non-negative integers m, n, s and N which satisfy the conditions in (1).*

Acknowledgment. I thank Professor Robert V. Moody for his help and support.

References

[1] Morikuni Goto & Frank D. Grosshans, *Semisimple Lie Algebras*, Lectures Notes in Pure and Applied Mathematics. New York and Basel: Marcel Dekker, Inc., 1978

[2] Nathan Jacobson, *Lie Algebras*. New York: Dover, 1962

[3] Robert V. Moody & Arturo Pianzola, *Lie Algebras with Triangular Decompositions*, Canadian Mathematical Society Series of Monographs and Advanced Texts. New York: Wiley-Inerscience, 1995